



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2350208 A1 2000/05/18

(21) **2 350 208**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 1999/11/09
(87) Date publication PCT/PCT Publication Date: 2000/05/18
(85) Entrée phase nationale/National Entry: 2001/05/08
(86) N° demande PCT/PCT Application No.: EP 99/08478
(87) N° publication PCT/PCT Publication No.: WO 00/27819
(30) Priorités/Priorities: 1998/11/10 (9824579.8) GB;
1999/03/03 (199 10 396.8) DE

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 213/38, A61K 31/4409,
A61P 35/00, A61P 17/00, A61K 31/166, C07D 265/26,
C07D 401/14, C07D 417/12, C07D 413/12,
C07D 409/12, C07D 405/12, C07D 401/12,
C07D 213/61, C07D 213/40, C07C 237/30

(71) Demandeurs/Applicants:
SCHERING AKTIENGESELLSCHAFT, DE;
NOVARTIS AKTIENGESELLSCHAFT, CH

(72) Inventeurs/Inventors:
SCHIRNER, MICHAEL, DE;
MENRAD, ANDREAS, DE;
OTTOW, ECKHARD, DE;
...

(74) Agent: MARKS & CLERK

(54) Titre : AMIDES D'ACIDE ANTHRANILIQUE ET LEUR UTILISATION COMME MEDICAMENT
(54) Title: ANTHRANILIC ACID AMIDES AND THE USE THEREOF AS MEDICAMENTS

(57) Abrégé/Abstract:

The invention relates to anthranilic acid amides and the use thereof as medicaments for the treatment of diseases that are triggered by persistent angiogenesis, in addition to intermediate products in the production of anthranilic acid amides.

(72) Inventeurs(suite)/Inventors(continued): KRUGER, MARTIN, DE; SEIDELMANN, DIETER, DE; HUTH, ANDREAS, DE; FERRARI, STEFANO, CH; BRUGGEN, JOSE, CH; MESTAN, JURGEN, DE; WOOD, JEANETTE MARJORIE, CH; FURET, PASCAL, FR; MANLEY, PAUL WILLIAM, CH; BOLD, GUIDO, CH; THIERAUCH, KARL-HEINZ, DE

ABSTRACT

The invention relates to anthranilic acid amides and the use thereof as medicaments for the treatment of diseases that are triggered by persistent angiogenesis, in addition to intermediate products in the production of anthranilic acid amides.

11/3/2000

EP 009908478


European Patent Office

GD1- Berlin Office

November 3, 2000

Claims

1. Compounds of general formula I



in which


A

stands for the group $=NR^2$,

W


stands for oxygen, sulfur, two hydrogen atoms or the group $=NR^8$,

Z

stands for the group $=NR^{10}$, $=N-$, or $-N(R^{10})-(CH_2)_q-$, branched or unbranched C_{1-6} alkyl or the group

AMENDED SHEET

or A, Z and R' together form the group



m, n and o

stand for 0-3,

q

stands for 1-6,

$R_a, R_b, R_c, R_d, R_e, R_f$, independently of one another, stand

for hydrogen, C₁₋₄ alkyl or the group

=NR¹⁰, and/or R_a and/or R_b can form a

bond with R_c and/or R_d or R_c can form a

bond with R_e and/or R_f ,

x

stands for the group $=NR^9$,

Y

stands for the group $-(CH_2)_p$,

p

stands for 1-4,

 \mathbb{R}^1

stands for C₁₋₆ alkyl that is

unsubstituted or is optionally

AMENDED SHEET


substituted in one or more places with halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, hydroxy, nitro, cyano or substituted in one or more places with halogen, or C₁₋₆-alkoxy-substituted naphthyl, biphenyl, phenyl, thiophenyl, furanyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, triazinyl, quinolinyl or isoquinolinyl; or 5-chloro-2,3-dihydroindeny, 2,3-dihydroindeny, thienyl, 6-fluoro-1H-indol-3-yl, 1,2,3,4-tetrahydronaphthyl, benzo-1,2,5-oxadiazole, 6,7-dimethoxy-1,2,3,4-tetrahydro-2-naphthyl, or for the group

AMENDED SHEET


AMENDED SHEET

whereby phenyl, substituted phenyl or naphthyl is not right in the $=NR^2$ group in the meaning of A,

R^2 stands for hydrogen or C_{1-6} alkyl,
 R^3 stands for naphthyl, biphenyl, phenyl, thiophenyl, furanyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, triazinyl, quinolinyll or isoquinolinyll that is unsubstituted or optionally substituted in one or more places with halogen; C_{1-6} alkyl, C_{1-6} alkoxy or hydroxy, or for the group



R^4 , R^5 , R^6 , and R^7 , independently of one another, stand for hydrogen, halogen, or C_{1-6} alkoxy, C_{1-6} alkyl or C_{1-6} carboxylalkyl that is unsubstituted or optionally substituted in one or more places with halogen,
 or R^5 and R^6 together form the group



R^8 , R^9 , and R^{10} , independently of one another, stand for hydrogen or C_{1-6} alkyl,

AMENDED SHEET

as well as their isomers and salts,

whereby


- a) If R^9 stands for hydrogen, Y stands for the group $-(CH_2)_p$, in which $p = 1$, R^3 stands for phenyl and W stands for oxygen, R^1 must not stand for unsubstituted imidazol-1-yl, and
- b) If W stands for oxygen, R^2 stands for hydrogen, R^9 stands for hydrogen,
- Y stands for the group $-(CH_2)_p-$, in which $p = 1-4$ and R^1 stands for pyridyl,

or

Y stands for the group $-(CH_2)_p-$, in which $p = 1$, and R^1 stands for phenyl or phenyl that is substituted in one place by methyl, chlorine or bromine, and Z stands for methyl or ethyl, R^3 must not stand for pyridyl, phenyl or phenyl that is substituted in one place by methyl, chlorine or bromine.


2. Compounds of general formula I, according to claim 1, in which

- A stands for the group $=NR^2$,
- W stands for oxygen, sulfur, two hydrogen atoms or the group $=NR^8$,
- Z stands for the group $=NR^{10}$, $=N-$ or $-N(R^{10})-(CH_2)_q-$, branched or unbranched C_{1-6} alkyl or the group



AMENDED SHEET


or A, Z and R¹ together form the group




- m, n, and o stand for 0-3,
q stands for 1-6,
R_a, R_b, R_c, R_d, R_e and R_f, independently of one another, stand for hydrogen, C₁₋₄ alkyl or the group =NR¹⁰,
X stands for the group =NR⁹,
Y stands for the group -(CH₂)_p,
p stands for 1-4,
R¹ stands for phenyl, pyridyl, 5-chloro-2,3-dihydroindenyl, 2,3-dihydroindenyl, thienyl, 6-fluoro-1H-indol-3-yl, naphthyl, 1,2,3,4-tetrahydronaphthyl, benzo-1,2,5-oxadiazole, 6,7-dimethoxy-1,2,3,4-tetrahydro-2-naphthyl or for phenyl or pyridyl that is substituted in one or more places with C₁-C₄ alkyl, C₁-

AMENDED SHEET

C₄, alkoxy, hydroxy, halogen, or
trifluoromethyl, or for the group



AMENDED SHEET




whereby phenyl, substituted phenyl or naphthyl is not right in the $=NR^2$ group in the meaning of A

R^2

stands for hydrogen or C_{1-6} alkyl,


R^3

stands for naphthyl, biphenyl, phenyl, thiophenyl, furanyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, triazinyl, quinolinyl or isoquinolinyl that is unsubstituted or that is optionally substituted in one or more places with halogen, C_{1-6} alkyl, C_{1-6} alkoxy or hydroxy, or for the group



AMENDED SHEET

R^4 , R^5 , R^6 and R^7 , independently of one another, stand for hydrogen, halogen or C_{1-6} alkoxy or C_{1-6} alkyl that is unsubstituted or optionally substituted in one or more places with halogen, or R^5 and R^6 together form the group




R^8 , R^9 and R^{10} , independently of one another, stand for hydrogen or C_{1-6} alkyl, as well as their isomers and salts, whereby, if W stands for oxygen, R^2 stands for hydrogen, R^9 stands for hydrogen, Y stands for the group $-(CH_2)_p$, in which $p = 1-4$, and R^1 stands for pyridyl, or

Y stands for the group $-(CH_2)_p$, in which $p = 1$, and R^1 stands for phenyl or phenyl that is substituted in one place by methyl, chlorine or bromine, and Z stands for methyl or ethyl, R^3 must not stand for pyridyl, phenyl or phenyl that is substituted in one place by methyl, chlorine or bromine.


3. Compounds of general formula I according to claims 1 and 2, in which

A	stands for the group $=NR^2$,
W	stands for oxygen, sulfur or two hydrogen atoms,
Z	stands for the group $=NR^{10}$, $=N-$, or $-N(R^{10})-(CH_2)_q-$ or the group

AMENDED SHEET




or A, Z and R¹ together form the group



- m, n and o stand for 0-3,
q stands for 1-6,
R_a, R_b, R_c, R_d, R_e, R_f, independently of one another, stand for hydrogen or methyl or the group =NR¹⁰,
X stands for the group =NR⁹,
Y stands for the group -CH₂-,
R¹ stands for phenyl, pyridyl, p-chlorophenyl, p-methylphenyl, p-methoxyphenyl, 5-chloro-2,3-dihydroindenyl, 2,3-dihydroindenyl,

AMENDED SHEET

thienyl, 6-fluoro-1H-indol-3-yl,
 naphthyl, 1,2,3,4-tetrahydronaphthyl,
 benzo-1,2,5-oxadiazole, 6,7-dimethoxy-
 1,2,3,4-tetrahydro-2-naphthyl, or for
 phenyl or pyridyl that is substituted in
 one or more places with C₁-C₄ alkyl, C₁-
 C₄ alkoxy, hydroxy, halogen, or
 trifluoromethyl, or for the group




AMENDED SHEET

whereby phenyl, or substituted phenyl or naphthyl is not right in the $=NR^2$ group in the meaning of A,
 R^2 stands for hydrogen or methyl,

AMENDED SHEET

R^3

stands for pyridyl or for phenyl or 1,2,3,4-tetrahydronaphthyl that is substituted in one or more places with hydroxy, halogen, methyl or methoxy, or for the group

 R^5 and R^6 ,

independently of one another, stand for hydrogen, halogen, methyl, methoxy or trifluoromethyl,

 R^4 and R^7 ,

independently of one another, stand for hydrogen,

 R^9

stands for hydrogen,

AMENDED SHEET

R^{10} stands for hydrogen or methyl,
as well as their isomers and salts, whereby if W stands for
oxygen, R^2 stands for hydrogen, R^9 stands for hydrogen,


R^1 stands for pyridyl,

or

R^1 stands for phenyl or phenyl that is substituted in one
place by methyl, chlorine or bromine, and Z stands for methyl or
ethyl, R^3 must not stand for pyridyl or phenyl that is
substituted in one place by methyl, chlorine or bromine.


4. Compounds of general formula I according to claims 1 to
3, in which

A stands for the group $=NR^2$,
W stands for oxygen,
Z stands for the group $=NR^{10}$, $=N-$, or
 $-N(R^{10})-(CH_2)_q-$ or the group



or A, Z and R^1 together form the group

AMENDED SHEET



m, n and o stand for 0-3,

q stands for 1-6,





































$R_a, R_b, R_c, R_d, R_e, R_f$, independently of one another, stand for hydrogen or methyl or the group $=NR^{10}$,

X stands for the group $=NR^9$,

Y stands for the group $-CH_2-$,

R^1 stands for phenyl, pyridyl, 5-chloro-2,3-dihydroindenyl, 2,3-dihydroindenyl, thienyl, 6-fluoro-1H-indol-3-yl, naphthyl, 1,2,3,4-tetrahydronaphthyl, benzo-1,2,5-oxadiazole or 6,7-dimethoxy-1,2,3,4-tetrahydro-2-naphthyl or for a phenyl or pyridyl that is substituted in one more places with C_1-C_4 alkyl, C_1-C_4 alkoxy, hydroxy, halogen, or trifluoromethyl, or for the group

AMENDED SHEET

AMENDED SHEET

whereby phenyl, or substituted phenyl or naphthyl is not right in the $=NR^2$ group in the meaning of A,

R^2

stands for hydrogen or methyl,

R^3


stands for pyridyl or for phenyl,

pyridyl or 1,2,3,4-tetrahydronaphthyl

that is substituted in one or more

places with hydroxy, halogen, methyl or

methoxy, or for the group



AMENDED SHEET

R⁵ and R⁶, independently of one another, stand for hydrogen, halogen, methyl, methoxy, or trifluoromethyl,

R⁴ and R⁷, independently of one another, stand for hydrogen and halogen,

R⁹ stands for hydrogen,

R¹⁰ stands for hydrogen or methyl,

as well as their isomers and salts, whereby if R² stands for hydrogen, R¹ stands for pyridyl,

or

R¹ stands for phenyl or phenyl that is substituted in one place by methyl, chlorine or bromine, and Z stands for methyl or ethyl, R³ must not stand for pyridyl or phenyl that is substituted in one place by methyl, chlorine or bromine.

5. Compounds of general formula I according to claims 1 to 3, in which


A stands for the group =NR²,

W stands for sulfur,


AMENDED SHEET

Z

stands for the group $=NR^{10}$, $=N-$, $-N(R^{10})-$
 $(CH_2)_q-$ or the group



or A, Z and R^1 together form the group



m, n and o

stand for 0-3,

q

stands for 1-6,

R_a , R_b , R_c , R_d , R_e , R_f , independently of one another, stand
 for hydrogen or methyl or the group
 $=NR^{10}$,

X

stands for the group $=NR^9$,



















Y

stands for the group $-CH_2-$,



















AMENDED SHEET

R¹

stands for phenyl, pyridyl, 5-chloro-2,3-dihydroindenyl, 2,3-dihydroindenyl, thienyl, 6-fluoro-1H-indol-3-yl, naphthyl, 1,2,3,4-tetrahydronaphthyl, benzo-1,2,5-oxadiazole or 6,7-dimethoxy-1,2,3,4-tetrahydro-2-naphthyl or for phenyl or pyridyl that is substituted in one or more places with C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, halogen, or trifluoromethyl, or for the group

AMENDED SHEET

whereby phenyl, or substituted phenyl or naphthyl is not right in the =NR² group in the meaning of A,


R²

stands for hydrogen or methyl,

R³

stands for pyridyl or for phenyl, pyridyl or 1,2,3,4-tetrahydronaphthyl that is substituted in one or more places with hydroxy, halogen, methyl or methoxy, or for the group

AMENDED SHEET



R^5 and R^6 ,

independently of one another, stand for hydrogen, halogen, methyl, methoxy or trifluoromethyl,

R^6 and R^7 ,

independently of one another, stand for hydrogen and halogen,

R^9

stands for hydrogen,

R^{10}


stands for hydrogen or methyl,

AMENDED SHEET


as well as their isomers and salts.

6. Compounds of general formula I according to claims 1 to 3, in which

A stands for the group $=NR^2$,
 W stands for two hydrogen atoms,
 Z stands for the group $=NR^{10}$, $=N-$, or
 $-N(R^{10})-(CH_2)_q-$ or the group



or A, Z, and R^1 together form the group



m, n and o

stand for 0-3,

q

stands for 1-6,


AMENDED SHEET

$R_a, R_b, R_c, R_d, R_e, R_f$, independently of one another, stand for hydrogen or methyl or the group $=NR^{10}$,


































X stands for the group $=NR^9$,

Y stands for the group $-CH_2-$,

R^1 stands for phenyl, pyridyl, 5-chloro-2,3-dihydroindenyl, 2,3-dihydroindenyl, thienyl, 6-fluoro-1H-indol-3-yl, naphthyl, 1,2,3,4-tetrahydronaphthyl, benzo-1,2,5-oxadiazole or 6,7-dimethoxy-1,2,3,4-tetrahydro-2-naphthyl or for a phenyl or pyridyl that is substituted in one or more places with C_1-C_4 alkyl, C_1-C_4 alkoxy, hydroxy, halogen, or trifluoromethyl, or for the group



AMENDED SHEET

AMENDED SHEET

whereby phenyl, or substituted phenyl or naphthyl is not right in the $=NR^2$ group in the meaning of A,

R^2

stands for hydrogen or methyl,

R^3


stands for pyridyl or for phenyl,

pyridyl or 1,2,3,4-tetrahydronaphthyl

that is substituted in one or more

places with hydroxy, halogen, methyl or


methoxy, or for the group



AMENDED SHEET

R^4 and R^7 , independently of one another, stand for hydrogen, halogen, methyl, methoxy or trifluoromethyl,
 R^5 and R^6 , independently of one another, stand for hydrogen and halogen,
 R^9 stands for hydrogen,
 R^{10} stands for hydrogen or methyl,
 as well as their isomers and salts.


7. Use of the compounds of general formula I




in which

A stands for the group $=NR^2$,
 W stands for oxygen, sulfur, two hydrogen atoms or the group $=NR^8$,
 Z stands for the group $=NR^{10}$, $=N-$ or $-N(R^{10})-(CH_2)_q-$, branched or unbranched C_{1-6} alkyl or the group

AMENDED SHEET



or A, Z and R¹ together form the group



m, n, and o stand for 0-3,

q stands for 1-6,


































R_a, R_b, R_c, R_d, R_e and R_f, independently of one another, stand for hydrogen, C₁₋₄ alkyl or the group =NR¹⁰, and/or R_a and/or R_b can form a bond with R_c and/or R_d or R_c can form a bond with R_e and/or R_f,

X stands for the group =NR⁹,

AMENDED SHEET

Y stands for the group $-(CH_2)_p$,
 p stands for 1-4,
 R¹ stands for C₁₋₆ alkyl that is
 unsubstituted or optionally substituted
 in one or more places with halogen, C₁₋₆
 alkyl, C₁₋₄ alkoxy, hydroxy, nitro,
 cyano, or in one or more places with
 halogen, or C₁₋₆ alkoxy-substituted
 naphthyl, biphenyl, phenyl, thiophenyl,
 furanyl, oxazolyl, thiazolyl,
 imidazolyl, pyrazolyl, pyridyl,
 pyrimidinyl, triazinyl, quinolinyl or
 isoquinolinyl; or 5-chloro-2,3-
 dihydroindenyl, 2,3-dihydroindenyl,
 thienyl, 6-fluoro-1H-indol-3-yl,
 1,2,3,4-tetrahydronaphthyl, benzo-1,2,5-
 oxadiazole, 6,7-dimethoxy-1,2,3,4-
 tetrahydro-2-naphthyl, or for the group

AMENDED SHEET

AMENDED SHEET


whereby phenyl, substituted phenyl or naphthyl is not right in the $=NR^2$ group in the meaning of A,

R^2

stands for hydrogen or C_{1-6} alkyl,


R^3

stands for naphthyl, biphenyl, phenyl, thiophenyl, furanyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, triazinyl, quinolinyl or isoquinolinyl that is unsubstituted or that is optionally substituted in one or more places with halogen, C_{1-6} alkyl, C_{1-6} alkoxy or hydroxy, or for the group



R^4 , R^5 , R^6 and R^7 ,

independently of one another, stand for hydrogen, halogen or C_{1-6} alkoxy, C_{1-6} alkyl or C_{1-6} carboxylalkyl that is unsubstituted or optionally substituted in one or more places with halogen, or R^5 and R^6 together form the group



AMENDED SHEET

R⁸, R⁹ and R¹⁰, independently of one another, stand for hydrogen or C₁₋₆ alkyl,

as well as their isomers and salts, for the production of a pharmaceutical agent for the treatment of tumors, psoriasis, arthritis, such as rheumatoid arthritis, hemangioma, angiofibroma, eye diseases, such as diabetic retinopathy, neovascular glaucoma, renal diseases, such as glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombic microangiopathic syndrome, transplant rejections and glomerulopathy, fibrotic diseases, such as cirrhosis of the liver, mesangial-cell-proliferative diseases, arteriosclerosis, injuries to the nerve tissue, and for inhibiting the reocclusion of vessels after balloon catheter treatment, in vascular prosthetics or after mechanical devices are used to keep vessels open, such as, e.g., stents.

8. Pharmaceutical agent that contains at least one compound according to claims 1 to 6.

9. Pharmaceutical agent according to claim 8 for the treatment of tumors, psoriasis, arthritis, such as rheumatoid arthritis, hemangioma, angiofibroma, eye diseases, such as diabetic retinopathy, neovascular glaucoma, renal diseases, such as glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombic microangiopathic syndrome, transplant rejections and glomerulopathy, fibrotic diseases, such as cirrhosis of the liver, mesangial-cell-proliferative diseases, arteriosclerosis, injuries to the nerve tissue, and for inhibiting the reocclusion of vessels after balloon catheter

AMENDED SHEET

treatment, in vascular prosthetics or after mechanical devices are used to keep vessels open, such as, e.g., stents.

10. Compounds according to claims 1 to 6 and pharmaceutical agents according to claims 6 and 8 with suitable formulations and vehicles.

11. Use of the compounds of formula I according to claims 1 to 7 as inhibitors of tyrosine kinases KDR and FLT.

12. Use of the compounds of general formula I according to claims 1 to 6 in the form of a pharmaceutical preparation for enteral, parenteral and oral administration.

AMENDED SHEET